

Research



Cite this article: Philson CS, Blumstein DT. 2023 Emergent social structure is typically not associated with survival in a facultatively social mammal. *Biol. Lett.* **19**: 20220511. <https://doi.org/10.1098/rsbl.2022.0511>

Received: 9 November 2022

Accepted: 23 February 2023

Subject Areas:

behaviour, evolution

Keywords:

behavioural ecology, social structure, fitness consequences, social network analysis, social selection, survival

Author for correspondence:

Conner S. Philson

e-mail: cphilson@ucla.edu

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6466944>.

Animal behaviour

Emergent social structure is typically not associated with survival in a facultatively social mammal

Conner S. Philson^{1,2} and Daniel T. Blumstein^{1,2}

¹Department of Ecology and Evolutionary Biology, University of California, 621 Young Drive South, Los Angeles, CA 90095-1606, USA

²Rocky Mountain Biological Laboratory, Box 519, Crested Butte, CO 81224, USA

CSP, 0000-0002-5974-347X; DTB, 0000-0001-5793-9244

For social animals, group social structure has important consequences for disease and information spread. While prior studies showed individual connectedness within a group has fitness consequences, less is known about the fitness consequences of group social structure for the individuals who comprise the group. Using a long-term dataset on a wild population of facultatively social yellow-bellied marmots (*Marmota flaviventris*), we showed social structure had largely no relationship with survival, suggesting consequences of individual social phenotypes may not scale to the group social phenotype. An observed relationship for winter survival suggests a potentially contrasting direction of selection between the group and previous research on the individual level; less social individuals, but individuals in more social groups experience greater winter survival. This work provides valuable insights into evolutionary implications across social phenotypic scales.

1. Introduction

An individual's social phenotype has important implications for its fitness in group-living species [1]. Survival, an important fitness correlate, is influenced by the degree, rate and context of social interactions in some species, as measured by social network analysis. For example, stronger social relationships have been associated with greater survival in male bottlenose dolphins (*Tursiops* sp.; [2]), chacma baboons (*Papio hamadryas ursinus*; [3]), Barbary macaques (*Macaca sylvanus*; [4,5]) and feral horses (*Equus caballus*; [6,7]). Several mechanisms contribute to this positive sociality–survival relationship, such as predator defence and detection [8–10] and resource exploitation [11,12]. Yet, higher rates of sociality can also be costly in terms of survival, as seen in female bottlenose dolphins [2] and is often associated with predator attraction and disease acquisition [13–15].

This previous work exclusively explored the fitness consequences of an individual's social phenotype, leaving the consequences of the group social phenotype unstudied [16–18]. Group social structure—emergent network traits generated by the interactions of all group members—quantifies the group social phenotype in a way more complex than averages of individual-level network values [19–22]. Social structure influences key ecological and evolutionary processes [23] such as group formation and regulation [24] and movement patterns [25]. Social structure is in turn influenced by ecological conditions [26,27] and group composition of individual social and non-social phenotypes [28,29]. In most species, social structures are non-random, repeatable across years, group compositions, and environmental gradients and conditions [18,29–37].

Social structure may also have fitness consequences. Captive bluebanded gobies (*Lythrypnus dalli*), an obligately social species, have reduced reproductive success when living in groups with more reciprocated aggressive interactions

Table 1. Measures of social structure with the *a priori* hypothesized direction of the relationship.

measure	description	references	interpretation	summer hypothesis	winter hypothesis
density	number of interactions observed represented as a fraction of all possible interactions	Burt [46]; Wasserman & Faust [20]; Grund [47]	how connected a group is	+	–
transitivity	proportion of completely connected triads out of the total possible triads	Wasserman & Faust [20]; Milo <i>et al.</i> [48]; Faust [49]	how cyclically connected a group is. There are more transitive components in affiliative networks in this system [50]	+	–
reciprocity	the number of mutual interactions divided by the number of possible mutual interactions	Wasserman & Faust [20]; Kankanhalli <i>et al.</i> [51]; Squartini <i>et al.</i> [52]	used to quantify how mutual or one-sided interactions are in a group	+	+
degree assortativity	tendency for social ties to share similar individual degree measures	McPherson <i>et al.</i> [53]; Currarini <i>et al.</i> [54]	how socially homogeneous a group is, in terms of individual's number of social partners	+	+
average path length	average of the shortest paths between all pairs of nodes	Watts & Strogatz [55]; Broder <i>et al.</i> [56]	identifies the size of a network, in addition to raw group size	–	+
cut points	number of social ties that if cut will result in two or more separate networks	Wasserman & Faust [20]; Borgatti [57]	how stable or fragmentable (breakable) a group is	–	+

[38]. In wild yellow-bellied marmots, adult female reproductive success modestly increases when living in more fragmentable social groups [39]. Adults also gain body mass more slowly when living in more connected groups, though yearlings gain mass more quickly in socially homogeneous groups [40]. These studies highlight how social structure may have different fitness consequences across species, demographic roles and social systems [16,17]. However, how social structure specifically relates to patterns of survival in animal populations is not understood.

We explored the relationship between group social structure and survival over both the active summer season and winter hibernation in a long-studied population of yellow-bellied marmots. In this system, an individual's social phenotype predicts survival [41,42] and other fitness correlates (alarm call propensity: [43]; reproductive success: [44]; longevity: [45]). Informed by previous work on the individual and group levels in this system, we developed *a priori* hypotheses (table 1) for six network measures, capturing the connectedness, mutuality and homophily of a group, and both summer and winter survival. Females living in larger social groups and yearling females engaging in more social interactions experience enhanced summer survival [42]. Predation is the primary driver of summer mortality in this system [58], thus more social individuals, or those residing in more connected groups, may better detect and avoid predators [8–10]. We predict that residing in more connected, reciprocal and

socially homogeneous groups will be associated with increased summer survival. However, more social marmots are less likely to survive hibernation [41]. Since body condition is the primary driver of winter mortality, this may be a function of social interaction time costs [59] and/or individuals in groups having more costly periodic arousals during social hibernation [60,61]. Thus, we predict that residing in more connected, less reciprocal and socially heterogeneous groups will be associated with decreased winter survival, as was seen for mass gain and reproductive success [39,40].

2. Methods

(a) Study system

Yellow-bellied marmots are harem-polygynous, facultatively social ground-dwelling squirrels living in matrilineal colonies with one to two territorial males [62,63]. The population at the Rocky Mountain Biological Laboratory in Colorado (38°57'N, 106°59'W; ca 2900 m elevation) has been continuously studied since 1962. These marmots are active for five months annually (mid-April to mid-September). Following their winter hibernation, they mate soon after emergence with yearlings dispersing and new pups emerging in late-June to early-July. Annually, most males and about half of females disperse as yearlings, typically resulting in movement out of the study area [63]. We only explore adult summer and winter survival because yearling dispersal creates uncertainty about yearling survival.

(b) Data collection

We repeatedly trapped and observed marmots during their active season from 2002 to 2020. All individuals studied in our population have unique non-toxic dye marks on their dorsal pelage, allowing accurate identification of interacting individuals and precise estimates of survival. A detailed ethogram and behavioural observational methodology are outlined in Blumstein *et al.* [64]. The initiator, recipient, location, time and type of each interaction is recorded, with most interactions (79%) occurring between identified individuals. The remaining 21% interactions could not be identified because of interacting individuals' posture or visual obstructions. We excluded these interactions from our data, which should not significantly influence our estimates of social structure [65].

Only adults and yearlings are included in our social interaction data because only these cohorts were present in spring, when social interactions were most common. Pups were excluded because of their mid-season emergence and primary interaction with their mother and each other [66]. We eliminated transients by excluding individuals observed or trapped fewer than five times in a given year [41,43–45]. Colony sites are grouped into higher and lower elevation sites (five are at higher and seven are at lower elevation sites). Higher elevation sites are approximately 166 m higher and experience harsher weather conditions [36,67,68].

(c) Social network measures

Directed and weighted interaction matrices were constructed from affiliative interactions for each year (2002 to 2020) the package 'igraph' (version 1.3.5; [69]) for R 4.2.0 [70]. We focused on affiliative interactions (e.g. allogrooming, greeting and play) because they relate to summer and winter survival on the individual level [41,42] and affiliative interactions comprised 88% of interactions. These affiliative matrices consisted of 38 968 social interactions between 726 individuals (626 of whom were observed across multiple years). A social group is defined as a network component (set of connected individuals with no other external connections) appearing naturally within a valley location (higher elevation or lower elevation) in a given year. This operationalization produced 137 social groups with group sizes ranging from 3 to 58 individuals with a mean of 20.51 (s.e. = 0.52). Details on identical behavioural observation and network methods can be found in Philson & Blumstein [39].

Six social network measures were calculated for each group to quantify social structure (table 1). Density, transitivity, average path length and cut points quantify connectivity; reciprocity and degree assortativity quantify homophily. We selected these measures due to their importance in our system [39,40,50], other systems [18,27,38,71], and because these network measures have analogous measures on the individual level (e.g. density and degree; transitivity and clustering coefficient; [18]), aiding our understanding of consequences of social behaviour across social phenotypic scales [20]. The reliability of the social network measures is facilitated by our observations of marmot social groups across their entire active season (mean n observations per individual across years = 28.81, range of each year = 6.79–75.14) and low rate of unknown individuals involved in social interactions [65,72,73]. Because group size is associated with many marmot social network measures (e.g. density, cut points; [35,39,40]), we standardized each measure not already 'standardized' (i.e. degree assortativity and cut points) by dividing the network measures by group size [20,39,40].

(d) Data analysis

Summer survival was defined as individuals seen or trapped after 1 August or in the following years. Winter survival was

defined as individuals seen the following year/s. Survival data were paired with network measures from the current active season (for summer survival) or the active season before hibernation (for winter survival). We fitted two generalized linear mixed models (summer and winter) with a binomial distribution and a bobyqa optimizer with 20 000 iterations using 'lme4' [74–76]. Model assumptions and zero inflation were checked after fitting. The summer survival model included the six network measures, social group size, age, sex, June mass, valley location and a predation index as fixed effects. The winter survival model included the six network measures, group size, age, sex, August mass, valley location, and the date bare ground was first visible because of melting snowpack in the centre of our colony areas. Models had 559 observations consisting of 252 unique individuals in 91 social groups across 19 years.

We included the individual attributes age, sex, mass and location because survival is multi-causal and we wished to account for important attributes with known fitness implications [77–79]. Group size was included due to its relationship with fitness correlates in this system [35,44]. Predation index is a binary variable calculated by whether the number of predator observations at that colony was below or above the median number of predator observations across all colony areas in that year [80], providing a value relative to all other years [42]. Individual ID and year were included as random effects to account for annual environmental and demographic differences [36,81,82] and individuals observed over multiple years.

All continuous variables were standardized (mean-centred and divided by 1 s.d. using the 'scale' function in base R; [83]). Group size was log₁₀ transformed before scaling, but we employed no other transformations. We checked for multicollinearity between fixed effects and found each had a correlation coefficient of less than 0.8 [84,85], though both density and group size had a VIF of greater than 5 in both the summer and winter models (electronic supplementary material, table S1). Because group size has known relationships with fitness in this system [35,39,40,44], we removed density from both models and the interpretation of the results for the measures of social structure did not change (electronic supplementary material, table S2). Since this suggests our models were relatively robust, we report models that included both density and group size here.

Marginal and conditional R^2 values for each model and the semi-partial marginal and conditional R^2 that estimate variance explained by each fixed effect were calculated using 'partR2' [86,87]. We estimated 95% confidence intervals for our R^2 values using 100 parametric bootstrap iterations. Figures were generated with 'ggplot2' [88].

3. Results

Contrary to our *a priori* hypothesis, we found a statistically significant negative main effect of cut points on winter survival ($B = -0.359$; $p = 0.031$; s.e. = 0.167; figure 1; table 2), suggesting individuals residing in more fragmentable—breakable into two or more separate groups of two or more individuals—social groups experienced reduced winter survival. Overall, five of the six measures of social structure did not have a significant statistical relationship with winter survival, suggesting the relationship is modest. This model had a marginal R^2 value of 6.28% and a conditional R^2 value of 7.93%. Cut points explained 1.11% of the marginal semi-partial R^2 variance further suggesting the relationship is modest.

Rejecting our *a priori* hypotheses, we found no significant main effects of social structure in our summer survival model (figure 2; table 2), suggesting social structure does not play a primary role in summer survival. This model




Figure 1. Relationship between winter survival and social structure. The predictor variable is scaled. Darker points indicate overlaid data.

explained 7.39% of the marginal variance and 19.17% of the conditional variance.

4. Discussion

We found no strong and consistent evidence that social structure is related to summer survival in this system. However, because one measure of social structure (cut points) was related to winter survival, group structure may have a limited impact in specific contexts. Because group social structure exists on a larger, emergent phenotypic scale than an individual's direct social interactions [16,22,89], it is logical that the relationship between social structure and fitness may be quite modest because of its indirect nature, as seen in previous work [18,38,39].

Social structure not relating to summer survival is surprising given that more connected groups might excel at predator detection and/or avoidance [8–10,90,91]. However, the lack of a statistically significant relationship may result from the emergent, and hence indirect nature of social structure [18–22]. Since these facultatively social marmots experience mostly fitness costs from more social individual and group phenotypes [39–41,43–45], the consequences of more social individual phenotypes may not scale linearly to the group social phenotype, as seen in male forked fungus beetles [18]. That is, residing in a more connected group may not incur the same benefits as increased individual sociality. A marmot may have modest benefits at an individual level of being more socially connected with others, but still regulate their social interactions by residing in less connected groups. Because of the few

associations with winter survival, the strength of group-level selection is likely less than the strength of individual-level selection for survival in this system. However, targeted exploration of multi-level selection is required to specifically test this hypothesis.

Interestingly, the one statistically significant relationship we identified (between winter survival and cut points) may have interesting evolutionary implications and further suggests that the individual social phenotype may not scale to the group phenotype [18]. In this system, more socially connected individuals have decreased winter survival [41], suggesting selection acts against more social individual phenotypes. As we have shown here, individuals residing in less fragmentable (or more closely socially connected) groups have increased winter survival, suggesting selection acts towards more social group phenotypes. The mechanisms for this are entangled. Winter survival is closely associated with body mass [81]. Thus, more social individuals in more social groups may have less time to forage to develop energy reserves and may be more likely to socially hibernate, increasing the risk of costly torpor interruptions [60,61]. However, residing in a less fragmentable group, where individuals may be more likely to share space and hear conspecific alarm calls [26,27,30,92] may facilitate predator avoidance and allow for more time to gain mass. This potential explanation is muddled, but not dismissed, by the lack of an observed relationship for summer survival.

Importantly, residing in a less fragmentable group does not necessarily imply that individuals within that group are significantly more social [19–22]. This also does not imply more connected social structures are beneficial in this species

Table 2. Model estimates, standard error, *p*-value, marginal and conditional and semi-partial part R^2 . Italic *p*-values indicate $p < 0.05$.

	winter survival			summer survival				
	Est. \pm s.e.	<i>p</i> -value	marginal part R^2 (%)	conditional part R^2 (%)	Est. \pm s.e.	<i>p</i> -value	marginal part R^2 (%)	conditional part R^2 (%)
intercept	0.968 \pm 0.188	<0.001	6.28 (5.47–14.53)	7.93 (5.66–19.59)	1.91 \pm 0.449	<0.001	7.39 (3.04–18.84)	19.17 (7.46–94.49)
density	-0.229 \pm 0.346	0.508	0.02 (0–8.66)	1.66 (0–14.04)	0.848 \pm 0.601	0.158	0.88 (0–12.56)	12.66 (1.06–94.03)
transitivity	-0.163 \pm 0.233	0.485	0.17 (0–8.81)	1.82 (0–14.18)	0.021 \pm 0.312	0.947	0.02 (0–11.72)	11.8 (0.4–93.97)
reciprocity	-0.119 \pm 0.172	0.49	0.36 (0–8.98)	2.01 (0–14.34)	-0.26 \pm 0.225	0.247	0.95 (0–12.62)	12.72 (1.12–94.04)
degree assortativity	0.093 \pm 0.16	0.563	0.09 (0–8.73)	1.74 (0–14.11)	0.173 \pm 0.244	0.477	0 (0–11.69)	11.76 (0.38–93.97)
cut points	-0.359 \pm 0.166	0.031	1.11 (0.23–9.68)	2.75 (0.33–15)	-0.069 \pm 0.21	0.742	0 (0–11.6)	11.67 (0.34–93.96)
average path	0.125 \pm 0.239	0.602	0 (0–8.65)	1.65 (0–14.03)	0.589 \pm 0.322	0.067	0.73 (0–12.41)	12.51 (0.91–94.02)
length								
group size	-0.665 \pm 0.286	0.02	1.25 (0.37–9.81)	2.9 (0.48–15.13)	-0.466 \pm 0.421	0.269	0 (0–11.47)	11.54 (0.29–93.95)
valley location	0.087 \pm 0.131	0.508	0.19 (0–8.82)	1.84 (0–14.19)	0.283 \pm 0.219	0.196	0.88 (0–12.55)	12.66 (1.05–94.03)
sex	0.709 \pm 0.18	<0.001	3.24 (2.39–11.68)	4.88 (2.56–16.89)	0.699 \pm 0.223	0.002	1.89 (0.28–13.53)	13.67 (2.04–94.1)
age	-0.076 \pm 0.122	0.531	0.1 (0–8.74)	1.75 (0–14.11)	-0.137 \pm 0.227	0.546	0.44 (0–12.13)	12.22 (0.62–94)
August mass	0.525 \pm 0.172	0.002	1.94 (1.07–10.46)	3.59 (1.2–15.74)	0.62 \pm 0.31	0.045	0 (0–11.45)	11.51 (0.28–93.95)
date of snowmelt	-0.122 \pm 0.142	0.39	0.23 (0–8.86)	1.88 (0–14.23)	0.066 \pm 0.184	0.718	0 (0–11.64)	11.71 (0.36–93.96)




Figure 2. Relationship between summer survival and social structure. The predictor variable is scaled. Darker points indicate overlaid data.

[39,40], but rather that in specific contexts some benefits to increased sociality may emerge at different social scales. As mentioned, the strength and potential contrasting direction of selection between individual and group traits in this system requires further exploration. Opposing selection has been observed across species for non-social traits [71,93–98], though less research investigates selection acting on social behaviour across social scales [95,96,99]. Bluebanded gobies [38], chacma baboons (*Papio hamadryas ursinus*; [3]) and social bumblebees (*Bombus* sp.; [100,101]) experience fitness benefits across social scales. However, not all species may experience, or experience via the same mechanisms, aligned fitness consequences across social scales [18].

Understanding the evolution of social behaviour requires directly measuring the fitness consequences of natural behavioural variation [38]. While the ranging fitness consequences of individual social network position is well documented [2,16,102–106], growing research supports group-level social network traits also have a range of fitness consequences across species [18,38–40]. This work highlights evolutionary implications across social scales and our main finding is that group social phenotype has a complex relationship with fitness that does not necessarily scale. Our observed relationship may further suggest complexity via contrasting selection between the individual and group social

phenotypes on winter survival. However, because the relationship is modest and little research investigates selection acting on behaviour across social scales, future work using a multi-level selection approach is essential to better understand the adaptive value of sociality.

Data accessibility. Data and code to replicate analyses are available from OSF at <https://doi.org/10.17605/OSF.IO/T93E8> [107].

Data are provided in the electronic supplementary material [108]

Authors' contributions. C.S.P.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, project administration, visualization, writing—original draft and writing—review and editing; D.T.B.: conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, resources, supervision, writing—original draft and writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no conflicts of interest.

Funding. Research was supported by the National Geographic Society, UCLA, National Science Foundation (grant nos. I.D.B.R.-0754247, D.E.B.-1119660, 1557130, D.B.I. 0242960, 0731346 and 1226713), Animal Behaviour Society, American Society of Mammologists, and Rocky Mountain Biological Laboratory.

Acknowledgements. We thank the previous marmoteers who collected data, Julien Martin for managing the database, the staff of Rocky Mountain Biological Laboratory for support, the editor and four anonymous reviewers for their helpful efforts.

References

- Alexander RD. 1974 The evolution of social behavior. *Annu. Rev. Ecol. Syst.* **5**, 325–383. (doi:10.1146/annurev.es.05.110174.001545)
- Stanton MA, Mann J. 2012 Early social networks predict survival in wild bottlenose dolphins.

- PLoS ONE* **7**, 47508. (doi:10.1371/journal.pone.0047508)
3. Cheney DL, Silk JB, Seyfarth RM. 2016 Network connections, dyadic bonds and fitness in wild female baboons. *R. Soc. Open Sci.* **3**, 160255. (doi:10.1098/rsos.160255)
 4. McFarland R, Majolo B. 2013 Coping with the cold: predictors of survival in wild Barbary macaques, *Macaca sylvanus*. *Biol. Lett.* **9**, 20130428. (doi:10.1098/rsbl.2013.0428)
 5. Lehmann J, Majolo B, McFarland R. 2016 The effects of social network position on the survival of wild Barbary macaques, *Macaca Sylvanus*. *Behav. Ecol.* **27**, 20–28. (doi:10.1093/beheco/arv169)
 6. Cameron EZ, Setsaas TH, Linklater WL. 2009 Social bonds between unrelated females increase reproductive success in feral horses. *Proc. Natl Acad. Sci. USA* **106**, 13 850–13 853. (doi:10.1073/pnas.0900639106)
 7. Nuñez CM, Adelman JS, Rubenstein DI. 2015 Sociality increases juvenile survival after a catastrophic event in the feral horse (*Equus caballus*). *Behav. Ecol.* **26**, 138–147. (doi:10.1093/beheco/aru163)
 8. Hamilton WD. 1971 Geometry for the selfish herd. *J. Theor. Biol.* **31**, 295–311. (doi:10.1016/0022-5193(71)90189-5)
 9. Janson CH, Goldsmith ML. 1995 Predicting group size in primates: foraging costs and predation risks. *Behav. Ecol.* **6**, 326–336. (doi:10.1093/beheco/6.3.326)
 10. Clutton-Brock TH, Gaynor D, Mcllraith GM, Maccoll ADC, Kansky R, Chadwick P, Manser M, Skinner JD, Brotherton PNM. 1999 Predation, group size and mortality in a cooperative mongoose, *Suricata suricatta*. *J. Anim. Ecol.* **68**, 672–683. (doi:10.1046/j.1365-2656.1999.00317.x)
 11. Blundell GM, Ben-David M, Bowyer RT. 2002 Sociality in river otters: cooperative foraging or reproductive strategies? *Behav. Ecol.* **13**, 134–141. (doi:10.1093/beheco/13.1.134)
 12. Lancaster JR, Wilson R, Espinoza RE. 2006 Physiological benefits as precursors of sociality: why banded geckos band. *Anim. Behav.* **72**, 199–207. (doi:10.1016/j.anbehav.2006.01.010)
 13. Freeland WJ. 1976 Pathogens and the evolution of primate sociality. *Biotropica* **8**, 12–24. (doi:10.2307/2387816)
 14. Côté IM, Poulin R. 1995 Parasitism and group size in social animals: a meta-analysis. *Behav. Ecol.* **6**, 159–165. (doi:10.1093/beheco/6.2.159)
 15. Lucatelli J, Mariano-Neto E, Japyassú HF. 2021 Social interaction, and not group size, predicts parasite burden in mammals. *Evol. Ecol.* **35**, 115–130. (doi:10.1007/s10682-020-10086-6)
 16. Croft DP, Darden SK, Wey TW. 2016 Current directions in animal social networks. *Curr. Opin. Behav. Sci.* **12**, 52–58. (doi:10.1016/j.cobeha.2016.09.001)
 17. Fisher DN, McAdam AG. 2017 Social traits, social networks and evolutionary biology. *J. Evol. Biol.* **30**, 2088–2103. (doi:10.1111/jeb.13195)
 18. Costello RA, Cook PA, Brodie III ED, Formica VA. 2023 Multilevel selection on social network traits differs between sexes in experimental populations of forked fungus beetles. *Evolution* **77**, 289–303. (doi:10.1093/evolut/qpac012)
 19. Hinde RA. 1976 Interactions, relationships and social structure. *Man* **11**, 1–17. (doi:10.2307/2800384)
 20. Wasserman S, Faust K. 1994 *Social network analysis: methods and applications*, vol. **8**. Cambridge, MA: Cambridge University Press.
 21. Ilany A, Booms AS, Holekamp KE. 2015 Topological effects of network structure on long-term social network dynamics in a wild mammal. *Ecol. Lett.* **18**, 687–695. (doi:10.1111/ele.12447)
 22. Kappeler PM. 2019 A framework for studying social complexity. *Behav. Ecol. Sociobiol.* **73**, 1–14. (doi:10.1007/s00265-018-2601-8)
 23. Kurvers RHJM, Krause J, Croft DP, Wilson ADM, Wolf M. 2014 The evolutionary and ecological consequences of animal social networks: emerging issues. *Trends Ecol. Evol.* **29**, 326–335. (doi:10.1016/j.tree.2014.04.002)
 24. Balasubramaniam K, Beisner B, Vandeleest J, Atwill E, McCowan B. 2017 Social buffering and contact transmission: network connections have beneficial and detrimental effects on *Shigella* infection risk among captive rhesus macaques. *PeerJ* **4**, e2630. (doi:10.7717/peerj.2630)
 25. Dufour V, Sueur C, Whiten A, Buchanan-Smith HM. 2011 The impact of moving to a novel environment on social networks, activity and wellbeing in two new world primates. *Am. J. Primatol.* **73**, 802–811. (doi:10.1002/ajp.20943)
 26. Leu ST, Farine DR, Wey TW, Sih A, Bull CM. 2016 Environment modulates population social structure: experimental evidence from replicated social networks of wild lizards. *Anim. Behav.* **111**, 23–31. (doi:10.1016/j.anbehav.2015.10.001)
 27. Costello RA, Cook PA, Formica VA, Brodie III ED. 2022 Group and individual social network metrics are robust to changes in resource distribution in experimental populations of forked fungus beetles. *J. Anim. Ecol.* **00**, 1–13.
 28. Farine DR, Montiglio PO, Spiegel O. 2015 From individuals to groups and back: the evolutionary implications of group phenotypic composition. *Trends Ecol. Evol.* **30**, 609–621. (doi:10.1016/j.tree.2015.07.005)
 29. Cook PA, Baker OM, Costello RA, Formica VA, Brodie III ED. 2022 Group composition of individual personalities alters social network structure in experimental populations of forked fungus beetles. *Biol. Lett.* **18**, 20210509. (doi:10.1098/rsbl.2021.0509)
 30. Godfrey SS, Sih A, Bull CM. 2013 The response of a sleepy lizard social network to altered ecological conditions. *Anim. Behav.* **86**, 763–772. (doi:10.1016/j.anbehav.2013.07.016)
 31. Shizuka D, Chaine AS, Anderson J, Johnson O, Laursen IM, Lyon BE. 2014 Across-year social stability shapes network structure in wintering migrant sparrows. *Ecol. Lett.* **17**, 998–1007. (doi:10.1111/ele.12304)
 32. Wilson ADM, Krause S, James R, Croft DP, Ramnarine IW, Borner KK, Clement RJG, Krause J. 2014 Dynamic social networks in guppies (*Poecilia reticulata*). *Behav. Ecol. Sociobiol.* **68**, 915–925. (doi:10.1007/s00265-014-1704-0)
 33. Aplin LM *et al.* 2015 Consistent individual differences in the social phenotypes of wild great tits, *Parus major*. *Anim. Behav.* **108**, 117–127. (doi:10.1016/j.anbehav.2015.07.016)
 34. Wilson ADM, Krause S, Ramnarine IW, Borner KK, Clément RJG, Kurvers RHJM, Krause J. 2015 Social networks in changing environments. *Behav. Ecol. Sociobiol.* **69**, 1617–1629. (doi:10.1007/s00265-015-1973-2)
 35. Maldonado-Chaparro AA, Hubbard L, Blumstein DT. 2015 Group size affects social relationships in yellow-bellied marmots (*Marmota flaviventris*). *Behav. Ecol.* **26**, 909–915. (doi:10.1093/beheco/arv034)
 36. Maldonado-Chaparro AA, Martin JG, Armitage KB, Oli MK, Blumstein DT. 2015 Environmentally induced phenotypic variation in wild yellow-bellied marmots. *J. Mammal.* **96**, 269–278. (doi:10.1093/jmammal/gyu006)
 37. Blaszczyk MB. 2018 Consistency in social network position over changing environments in a seasonally breeding primate. *Behav. Ecol. Sociobiol.* **72**, 11. (doi:10.1007/s00265-017-2425-y)
 38. Solomon-Lane TK, Pradhan DS, Willis MC, Grober MS. 2015 Agonistic reciprocity is associated with reduced male reproductive success within harem social networks. *Proc. R. Soc. B* **282**, 20150914. (doi:10.1098/rspb.2015.0914)
 39. Philson CS, Blumstein DT. 2023 Group social structure has limited impact on reproductive success in a wild mammal. *Behav. Ecol.* **34**, 89–98. (doi:10.1093/beheco/ara102)
 40. Philson CS, Todorov S, Blumstein DT. 2022 Residing in socially connected and dissimilar groups relates to depressed mass gain rates for yearling marmots. *Behav. Ecol.* **33**, 115–125. (doi:10.1093/beheco/arab114)
 41. Yang WJ, Maldonado-Chaparro A, Blumstein DT. 2017 A cost of being amicable in a hibernating mammal. *Behav. Ecol.* **28**, 11–19. (doi:10.1093/beheco/arw125)
 42. Montero AP, Williams DM, Martin JGA, Blumstein DT. 2020 More social female yellow-bellied marmots, *Marmota flaviventris*, have enhanced summer survival. *Anim. Behav.* **160**, 113–119. (doi:10.1016/j.anbehav.2019.12.013)
 43. Fuong H, Maldonado-Chaparro A, Blumstein DT. 2015 Are social attributes associated with alarm calling propensity? *Behav. Ecol.* **26**, 587–592. (doi:10.1093/beheco/aru235)
 44. Wey TW, Blumstein DT. 2012 Social attributes and associated performance measures in marmots: bigger male bullies and weakly affiliating females have higher annual reproductive success. *Behav. Ecol. Sociobiol.* **66**, 1075–1085. (doi:10.1007/s00265-012-1358-8)
 45. Blumstein DT, Williams DM, Lim AN, Kroeger S, Martin JGA. 2018 Strong social relationships are

- associated with decreased longevity in a facultatively social mammal. *Proc. R. Soc. B* **285**, 20171934. (doi:10.1098/rspb.2017.1934)
46. Burt RS. 1992 Structural holes. Cambridge, UK: Cambridge University Press.
 47. Grund TU. 2012 Network structure and team performance: the case of English Premier League soccer teams. *Soc. Netw.* **34**, 682–690. (doi:10.1016/j.socnet.2012.08.004)
 48. Milo R, Shen-Orr S, Itzkovitz S, Kashtan N, Chklovskii D, Alon U. 2002 Network motifs: simple building blocks of complex networks. *Science* **298**, 824–827. (doi:10.1126/science.298.5594.824)
 49. Faust. 2010 A puzzle concerning triads in social networks: graph constraints and the triad census. *Soc. Netw.* **32**, 221–233. (doi:10.1016/j.socnet.2010.03.004)
 50. Wey TW, Jordán F, Blumstein DT. 2019 Transitivity and structural balance in marmot social networks. *Behav. Ecol. Sociobiol.* **73**, 88. (doi:10.1007/s00265-019-2699-3)
 51. Kankanhalli A, Tan BC, Wei KK. 2005 Contributing knowledge to electronic knowledge repositories: an empirical investigation. *MIS Q.* **29**, 113–143. (doi:10.2307/25148670)
 52. Squartini T, Picciolo F, Ruzzenenti F, Garlaschelli D. 2013 Reciprocity of weighted networks. *Sci. Rep.* **3**, 1–9. (doi:10.1038/srep02729)
 53. McPherson M, Smith-Lovin L, Cook JM. 2001 Birds of a feather: homophily in social networks. *Annu. Rev. Sociol.* **27**, 415–444. (doi:10.1146/annurev.soc.27.1.415)
 54. Currarini, Matheson J, Vega-Redondo F. 2016 A simple model of homophily in social networks. *Eur. Econ. Rev.* **90**, 18–19. (doi:10.1016/j.eurocorev.2016.03.011)
 55. Watts, Strogatz. 1998 Collective dynamics of ‘small-world’ networks. *Nature* **393**, 440–442. (doi:10.1038/30918)
 56. Broder A, Kumar R, Maghoul F, Raghavan P, Rajagopalan S, Stata R, Tomkins A, Wiener J. 2000 Graph structure in the web. *Comput. Netw.* **33**, 309–320. (doi:10.1016/S1389-1286(00)00083-9)
 57. Borgatti SP. 2006 Identifying sets of key players in a social network. *Comput. Math. Organ. Theory* **12**, 21–34. (doi:10.1007/s10588-006-7084-x)
 58. Van Vuren D. 2001 Predation on yellow-bellied marmots (*Marmota flaviventris*). *Am. Midl. Nat.* **145**, 94–100. (doi:10.1674/0003-0031(2001)145[0094:POYBMM]2.0.CO;2)
 59. Pollard KA, Blumstein DT. 2008 Time allocation and the evolution of group size. *Anim. Behav.* **76**, 1683–1699. (doi:10.1016/j.anbehav.2008.08.006)
 60. Arnold W. 1990 The evolution of marmot sociality: I. Why disperse late? *Behav. Ecol. Sociobiol.* **27**, 229–237.
 61. Armitage KB, Blumstein DT, Woods BC. 2003 Energetics of hibernating yellow-bellied marmots (*Marmota flaviventris*). *Comp. Biochem. Phys. A* **134**, 101–114. (doi:10.1016/S1095-6433(02)00219-2)
 62. Frase BA, Hoffmann RS. 1980 *Marmota flaviventris*. *Mamm. Species* **15**, 1–8. (doi:10.2307/3503965)
 63. Armitage KB. 1991 Social and population dynamics of yellow-bellied marmots: results from long-term research. *Annu. Rev. Ecol. Syst.* **22**, 379–407. (doi:10.1146/annurev.es.22.110191.002115)
 64. Blumstein DT, Wey TW, Tang K. 2009 A test of the social cohesion hypothesis: interactive female marmots remain at home. *Proc. R. Soc. B* **276**, 3007–3012. (doi:10.1098/rspb.2009.0703)
 65. Silk MJ, Jackson AL, Croft DP, Colhoun K, Bearhop S. 2015 The consequences of unidentifiable individuals for the analysis of an animal social network. *Anim. Behav.* **104**, 1–11. (doi:10.1016/j.anbehav.2015.03.005)
 66. Nowicki S, Armitage KB. 1979 Behavior of juvenile yellow-bellied marmots: play and social integration. *Z. Tierpsychol.* **51**, 85–105. (doi:10.1111/j.1439-0310.1979.tb00674.x)
 67. Van Vuren DH, Armitage KB. 1991 Duration of snow cover and its influence on life-history variation in yellow-bellied marmots. *Can. J. Zool.* **69**, 1755–1758. (doi:10.1139/z91-244)
 68. Blumstein DT, Ozgul A, Yovovitch V, Van Vuren DH, Armitage KB. 2006 Effect of predation risk on the presence and persistence of yellow-bellied marmot (*Marmota flaviventris*) colonies. *J. Zool.* **27**, 132–138. (doi:10.1111/j.1469-7998.2006.00098.x)
 69. Csardi G, Nepusz T. 2006 The igraph software package for complex network research. *Int. J. Commun. Syst.* **1695**, 1–9. version 1.2.11.
 70. R Development Core Team. 2021 *R: a language and environment for statistical computing*, version 4.1.2. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
 71. Weing C, Johnston JA, Willis CG, Maloof JN. 2007 Antagonistic multilevel selection on size and architecture in variable density settings. *Evolution* **61**, 58–67. (doi:10.1111/j.1558-5646.2007.00005.x)
 72. Davis GH, Crofoot MC, Farine DR. 2018 Estimating the robustness and uncertainty of animal social networks using different observational methods. *Anim. Behav.* **141**, 29–44. (doi:10.1016/j.anbehav.2018.04.012)
 73. Sánchez-Tójar A, Schroeder J, Farine DF. 2018 A practical guide for inferring reliable dominance hierarchies and estimating their uncertainty. *J. Anim. Ecol.* **87**, 594–608. (doi:10.1111/1365-2656.12776)
 74. Bates D, Maechler M, Nash MJ, Varadhan CR. 2014 minqa: Derivative-free optimization algorithms by quadratic approximation. R package version 1.2.4. See <http://CRAN.R-project.org/package=minqa>.
 75. Bates D, Maechler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
 76. Bates D, Maechler M, Bolker B, Walker S. 2015 lme4: Linear mixed-effects models using ‘Eigen’ and S4, R package version 1.1-27.1. See <http://CRAN.R-project.org/package=lme4> (accessed 20 July 2022).
 77. Wey TW, Blumstein DT. 2010 Social cohesion in yellow-bellied marmots is established through age and kin structuring. *Anim. Behav.* **79**, 1343–1352. (doi:10.1016/j.anbehav.2010.03.008)
 78. Huang B, Wey TW, Blumstein DT. 2011 Correlates and consequences of dominance in a social rodent. *Ethology* **117**, 573–585. (doi:10.1111/j.1439-0310.2011.01909.x)
 79. Blumstein DT, Keeley KN, Smith JE. 2016 Fitness and hormonal correlates of social and ecological stressors of female yellow-bellied marmots. *Anim. Behav.* **112**, 1–11. (doi:10.1016/j.anbehav.2015.11.002)
 80. Nash AL, Jebb AHM, Blumstein ET. 2020 Is the propensity to emit alarm calls associated with health status? *Curr. Zool.* **66**, 607–614. (doi:10.1093/cz/zoaa020)
 81. Kroeger SB, Blumstein DT, Armitage KB, Reid JM, Martin JG. 2018 Age, state, environment, and season dependence of senescence in body mass. *Ecol. Evol.* **8**, 2050–2061. (doi:10.1002/ece3.3787)
 82. Heissenberger S, de Pinho GM, Martin JG, Blumstein DT. 2020 Age and location influence the costs of compensatory and accelerated growth in a hibernating mammal. *Behav. Ecol.* **31**, 826–833. (doi:10.1093/beheco/araa013)
 83. Becker RA, Chambers JM, Wilks AR. 1988 *The new S language*. Belmont, CA: Wadsworth & Brooks/Cole.
 84. Franke GR. 2010 Multicollinearity. In *Wiley international encyclopedia of marketing* (eds J Sheth, N Malhotra). Chichester, UK: John Wiley & Sons, Ltd. (doi:10.1002/9781444316568.wiem02066)
 85. Shrestha N. 2020 Detecting multicollinearity in regression analysis. *Am. J. Appl. Math. Stat.* **8**, 39–42. (doi:10.12691/ajams-8-2-1)
 86. Nakagawa S, Schielzeth H. 2013 A general and simple method for obtaining R² from generalized linear mixed-effects models. *Methods Ecol. Evol.* **4**, 133–142. (doi:10.1111/j.2041-210x.2012.00261.x)
 87. Stoffel MA, Nakagawa S, Schielzeth H. 2021 partR2: Partitioning R² in generalized linear mixed models. *PeerJ* **9**, e11414. (doi:10.7717/peerj.11414)
 88. Wickham H. 2016 *Ggplot2: elegant graphics for data analysis*. Version 3.3.5. New York, NY: Springer-Verlag. See <https://ggplot2.tidyverse.org>
 89. Moore AJ, Brodie III ED, Wolf JB. 1997 Interacting phenotypes and the evolutionary process: I. Direct and indirect genetic effects of social interactions. *Evolution* **51**, 1352–1362. (doi:10.2307/2411187)
 90. Turner GF, Pitcher TJ. 1986 Attack abatement: a model for group protection by combined avoidance and dilution. *Am. Nat.* **128**, 228–240. (doi:10.1086/284556)
 91. Wrona FJ, Dixon RJ. 1991 Group size and predation risk: a field analysis of encounter and dilution effects. *Am. Nat.* **137**, 186–201. (doi:10.1086/285153)
 92. He P, Maldonado-Chaparro AA, Farine DR. 2019 The role of habitat configuration in shaping social structure: a gap in studies of animal social complexity. *Behav. Ecol. Sociobiol.* **73**, 1–14. (doi:10.1007/s00265-018-2602-7)
 93. Stevens L, Goodnight CJ, Kalisz S. 1995 Multilevel selection in natural populations of *Impatiens capensis*. *Am. Nat.* **145**, 513–526. (doi:10.1086/285753)

94. Tsuji K. 1995 Reproductive conflicts and levels of selection in the ant *Pristomyrmex pungens*: contextual analysis and partitioning of covariance. *Am. Nat.* **146**, 586–607. (doi:10.1086/285816)
95. Eldakar OT, Wilson DS, Dlugos MJ, Pepper JW. 2010 The role of multilevel selection in the evolution of sexual conflict in the water strider *Aquarius remigis*. *Evolution* **64**, 3183–3189. (doi:10.1111/j.1558-5646.2010.01087.x)
96. Laiolo P, Obeso JR. 2012 Multilevel selection and neighbourhood effects from individual to metapopulation in a wild passerine. *PLoS ONE* **7**, e38526. (doi:10.1371/journal.pone.0038526)
97. Van Vliet S, Doebeli M. 2019 The role of multilevel selection in host microbiome evolution. *Proc. Natl Acad. Sci. USA* **116**, 20 591–20 597. (doi:10.1073/pnas.1909790116)
98. Cordes LS, Blumstein DT, Armitage KB, CaraDonna PJ, Childs DZ, Gerber BD, Martin JGA, Oli MK, Ozgul A. 2020 Contrasting effects of climate change on seasonal survival of a hibernating mammal. *Proc. Natl Acad. Sci. USA* **117**, 18 119–18 126. (doi:10.1073/pnas.1918584117)
99. Royle NJ, Pike TW, Heeb P, Richner H, Kölliker M. 2012 Offspring social network structure predicts fitness in families. *Proc. R. Soc. B* **279**, 4914–4922. (doi:10.1098/rspb.2012.1701)
100. Otterstatter MC, Thomson JD. 2007 Contact networks and transmission of an intestinal pathogen in bumble bee (*Bombus impatiens*) colonies. *Oecologia* **154**, 411–421. (doi:10.1007/s00442-007-0834-8)
101. Figueroa LL, Grab H, Ng WH, Myers CR, Graystock P, McFrederick QS, McArt SH. 2020 Landscape simplification shapes pathogen prevalence in plant–pollinator networks. *Ecol. Lett.* **23**, 1212–1222. (doi:10.1111/ele.13521)
102. Wey TW, Blumstein DT, Shen W, Jordán F. 2008 Social network analysis of animal behavior: a promising tool for the study of sociality. *Anim. Behav.* **75**, 333–344. (doi:10.1016/j.anbehav.2007.06.020)
103. Whitehead H. 2008 *Analyzing animal societies: quantitative methods for vertebrate social analysis*. Chicago, IL: University of Chicago Press.
104. Formica VA, Wood CW, Larsen WB, Butterfield RE, Augat ME, Hougen HY, Brodie III ED. 2012 Fitness consequences of social network position in a wild population of forked fungus beetles (*Bolitotherus cornutus*). *J. Evol. Biol.* **25**, 130–137. (doi:10.1111/j.1420-9101.2011.02411.x)
105. Pinter-Wollman N *et al.* 2014 The dynamics of animal social networks: analytical, conceptual, and theoretical advances. *Behav. Ecol.* **25**, 242–255. (doi:10.1093/beheco/art047)
106. Farine DR, Sheldon BC. 2015 Selection for territory acquisition is modulated by social network structure in a wild songbird. *J. Evol. Biol.* **28**, 547–556. (doi:10.1111/jeb.12587)
107. Philson CS, Blumstein DT. 2022 Data from: Emergent social structure is typically not associated with survival in a facultatively social mammal. OSF (doi:10.17605/OSF.IO/T93E8)
108. Philson CS, Blumstein DT. 2023 Emergent social structure is typically not associated with survival in a facultatively social mammal. Figshare. (doi:10.6084/m9.figshare.c.6466944)